




DAMAANSA HOLDINGS

FAITH™ Drops



Faith

DOSAGE TABLE &
SUGGESTED TREATMENTS

ALL RIGHTS RESERVED

This manual remains the property of Damaansa Holdings Pty Ltd .
No part may be reproduced or copied without permission.

© Damaansa Holdings Pty Ltd
Head Office: PoBox 46816 • Glosderry • 7702
Tel: +27-21-6575786
Fax: +27-21-6575756
Cell: +27-(0)827746274

Regional Office: PoBox 95 • Witelsbos • 6304
Cell: +27. (0) 823399799
Fax +27. (0) 821313399799

(Associated Company: Anno-Organix Pty Ltd.).

WARNING: Any and all information and/or statements made within this document or on our website are for educational use only and are not intended to diagnose, treat or cure, prevent disease or replace the advice of a licensed Healthcare Practitioner.

Note also that our website may provide certain links to other websites operated by other parties. These links are provided for your convenience and reference only and we accept no liability or responsibility for their content.

Damaansa Holdings and its associated companies do not dispense medical advice, prescribe or diagnose illness. Any ideas or views expressed by Damaansa Holdings is not intended to be a substitute for conventional medical advice or service. By making use of our manual or product you agree that no responsibility or liability will be incurred to any person with respect to loss damage or injury caused or alleged to be caused directly or indirectly by the information contained in this document or on our website. Consult a licensed Healthcare Practitioner before beginning any alternative protocols.

RECOMMENDED TREATMENTS

This section of the document pertains specifically to research completed over more than 15 years.

The treatments indicated are a guideline only and are easily adjustable according to the patient's requirements.

Curing Diseases

We do not suggest that the FAITH™ Drops cures disease we say only that FAITH™ Drops supplies the immune system with the tools it needs to fight disease. The fact that the FAITH™ Drops do not attack healthy cells is evidence that the immune system is in control. When using the FAITH™ Drops the immune system becomes more efficient.

About the size of drops:

I refer in all cases to the standard FAITH™ bottle and cap shown in this book, which is a 30ml bottle with a dropper cap.

The 30ml bottle holds approximately 450 drops. One milliliter (or one CC) is equal to approximately 15 drops.



Fig.1. A standard FAITH™ 30ml HDPE bottle with dropper cap



Fig.2. A standard FAITH™ 30ml HDPE dropper cap

GENERAL INFORMATION :

When we refer to **FAITH™ Drops**, we refer to the bottle containing the 12 Herbal Tinctures / extracts in a Stabilised Chlorine Dioxide solution.

When we refer to **Activator**, we refer to the bottle containing the 9% Citric Acid solution.

When we refer to **Mixers**, we refer to any of the following:

BLACK Herbal tea, Rooibos tea, Coffee, Ceylon Tea. **OR**

FRUIT juices that DO NOT contain VITAMIN C - PREFERABLY, grape Juice, apple juice, pineapple juice or cranberry juice would be suitable provided they have no VITAMIN C added.

WHAT IS A PATHOGEN?

A pathogen is described as any biological agent that causes disease or illness to its host. Types of pathogens include Bacteria, Protozoa, Fungi, Parasites and Proteins.

Examples and/or typical effect of Bacteria Pathogens include:

Urinary Tract infections
Peritonitis
Typhoid
Foodborn illness such as Salmonella or E. coli
Tuberculosis
Anthrax
Toxic Shock Syndrome
Strep Throat
Pneumonia
Stomach Ulcers
Tularemia
Lyme Disease

Examples and/or typical effect of Virus Pathogens include:

HIV / Aids
Hepatitis A, B, C, D and E (liver disease)
Herpes
HPV (i.e. Genital Warts; HPV infection is a necessary factor in the development of nearly all cases of Cervical Cancer).
Warts
Chicken Pox, Small Pox, Cold Sores and Measles
Colds, Influenza Virus (i.e. H5N1, "Bird Flu")

Examples and/or typical effect of Protozoa Pathogens include:

Malaria
Giardiasis
Cryptosporidiosis
Chagas Disease

Examples and/or typical effect of Fungi Pathogens include:

Opportunistic Pneumonia
Ringworm
Candidiasis (Yeast Infection)
Histoplasmosis (i.e. "Darlings Disease")
Cryptococcosis

Examples and/or typical effect of Parasite Pathogens include:

Roundworm
Tapeworm

Examples and/or typical effect of Protein Pathogens include:

BSE (Mad Cow Disease)
vCJD (Creutzfeldt-Jakob Disease)

We quote below from excerpts from the **Research Paper “War Against Microbes” written by the Bradford Research Institute**, wherein it describes the virus inactivation as a result of a product called Dioxychlor®. This product is an inorganic sodium salt which under acidic conditions becomes unstable and decomposes into a variety of products including but not limited to chloride ions, hyperchlorous ions and nascent (atomic) oxygen., resulting in a neutral molecule consisting of three electronegative atoms held together by covalent and coordinate covalent bonds. From this cluster, a single atom of highly reactive nascent oxygen is liberated onto the target organisms. This product has the same active ingredient as FAITH Drops which is derived from activated Chlorine Dioxide and so this process is the process that Faith Drops undergoes when activated with citric acid. One possible mechanism for the liberation of nascent oxygen from the chlorite ion (ClO_2^-) in an acidic environment involves the association or binding of “H” (proton normally bound to the oxygen of water as H_2O) to any of the three pairs of unused electrons in the outer shell of chlorine.

Virus Inactivation:

A virus typically consists of an outer shell or coating of protein encapsulating a nucleic acid which may be either DNA or RNA (a retrovirus). The skeletal backbone of the nucleic acids includes derivatives of phosphoric acid (H_3PO_4 a very strong acid) in which two of the original three hydroxyl groups (-OH) are substituted, leaving only one active hydroxyl group per phosphate.

Some viruses may have some glycoproteins incorporated into their protein coats that is proteins to which polysaccharides (sugar chains) have been attached. The bound polysaccharides may attach to specific sites on the protein coat, effectively converting a protein surface into a polysugar surface. This new surface has specificity for certain polysaccharides found on the surface of specific cell types, thereby conveying specificity to virus binding as well as a degree of immunological protection.

Once bound to the appropriate cell type, the nucleic acid component of the virus is injected into the cell and in many ways, takes over the protein synthesis processes of the cell. Certain segments of the viral nucleic acid consist of genes that are responsible for the replication of the coat. The nucleic acid component replicates by a process known as “base pairing” in which each base of the original strands attracts and binds the corresponding base, forming a pair (A-T and C-G, described more fully below) through hydrogen bonding. The result is the replication of the complete virus until the cell bursts, releasing many additional viral particles into the surrounding medium. In the presence of these acidic nucleic acids, the dioxychlor molecule becomes unstable and releases nascent oxygen into the medium.

Nucleic acids, both RNA and DNA have many characteristics in common in fact, they are almost chemically identical. Each consists of a chain of alternating sugar (a modification of ribose, deoxyribose) and phosphate groups, known as the “phosphate – sugar backbone”. Attached to a specific carbon of each deoxyribose group is an organic ring compound known as a “base”. Altogether there are only 4 types of bases found in DNA namely, guanine “G”, cytosine “C”, one segment from DNA differ from another. RNA differs chemically from DNA in that the base thymine is replaced by the base uracil “U”, representing a very subtle bio-chemical difference. There is also a subtle difference in the sugar deoxyribose. .

The base guanine found in both RNA and DNA, is very sensitive to oxidation, forming 8-oxoguanine as the oxidation product. The release of Dioxychlor® results in the oxidation of the guanine residue with a formation of 8-oxoguanine, thereby disallowing the replication of the viral nucleic acid base pairing. Although the replication of the protein coat may continue, the formation of a complete functional virus has been blocked by Dioxychlor® oxidation.

For the full report go to the Dioxychlor® website

THE STANDARD FAITH™ FORMULA PROTOCOL

The standard FAITH™ formula is administered or taken a maximum of three times per day and each dose does NOT EXCEED more than 3 drops per dose and a maximum of 9 drops per day. We recommend that the last dose be taken at bed time. Reduce the number of drops immediately nausea is experienced. Increase number of drops one at a time until just under nausea level.

Day 1 - First dose:

- One (1) drop of FAITH™ Immune Fortifier in a clean glass.
- Add to these drops 1 drop of **ACTIVATOR**.
- Swirl the glass to mix the activator and FAITH™ drops together. (Without the **ACTIVATOR** it will not work).
- Set the timer and wait three (3) minutes.
- After three minutes, gently blow once or twice onto the surface of the mixture to remove the gas build up and then add to the activated FAITH™ ¼ of a glass of tap water or any suitable **MIXER**.

Drink this solution IMMEDIATELY.

If no nausea was experienced continue with the protocols as follows:

Day 1 - Second dose:

- Two (2) drops of FAITH™ Immune Fortifier in a clean glass.
- Add to these drops 2 drops of **ACTIVATOR**
- Continue with the protocols as for the first dose.

Drink this solution IMMEDIATELY.

If no nausea was experienced continue with the protocols as follows:

Day 1 - Third dose:

- Three (3) drops of FAITH™ Immune Fortifier.
- Add to these drops 3 drops of **ACTIVATOR**
- Continue with the protocols as for the previous doses

Drink this solution IMMEDIATELY.

If no nausea was experienced continue with the protocols as follows:

Continue with protocols as per Day 1. Do not exceed a total of nine (9) drops per day. Follow Standard Faith Formula Protocol for until patient has recovered.

THE AGGRESSIVE FAITH™ FORMULA PROTOCOL

The Aggressive FAITH™ formula is administered or taken a maximum of eight times per day (every hour) and each dose does NOT EXCEED more than 15 drops per dose. We recommend that the last dose be taken at bed time. If nausea is experienced reduce the number of drops immediately drop by drop until nausea recedes. Continue with protocols, slowly increasing the dosage drop by drop with each dose to just under nausea level.

Day 1 - First dose:

- Three (3) drops of FAITH™ Immune Fortifier in a clean glass.
- Add to these drops 3 drops of **ACTIVATOR**.
- Swirl the glass to mix the activator and FAITH™ drops together. (Without the **ACTIVATOR** it will not work).
- Set the timer and wait three (3) minutes.
- After three minutes, gently blow once or twice onto the surface of the mixture to remove the gas build up and then add to the activated FAITH™ ¼ of a glass of tap water or any suitable **MIXER**.

Drink this solution IMMEDIATELY.

If no nausea was experienced continue with the protocols as follows:

Day 1 - Second dose:

- Four (4) drops of FAITH™ Immune Fortifier in a clean glass.
- Add to these drops 4 drops of **ACTIVATOR**
- Continue with the protocols as for the first dose.

If no nausea was experienced continue with the protocols as follows:

Day 1 - Third dose:

- Five (5) drops of FAITH™ Immune Fortifier.
- Add to these drops 5 drops of **ACTIVATOR**
- Continue with the protocols as for the previous doses

If no nausea was experienced continue with the protocols as follows:

Day 1 - Fourth dose:

- Six (6) drops of FAITH™ Immune Fortifier.
- Add to these drops 6 drops of **ACTIVATOR**
- Continue with the protocols as for the previous doses

If no nausea was experienced continue with the protocols as follows:

Day 1 - Fifth dose:

- Seven (7) drops of FAITH™ Immune Fortifier.
- Add to these drops 7 drops of **ACTIVATOR**
- Continue with the protocols as for the previous doses

If no nausea was experienced continue with the protocols as follows:

Day 1 - Sixth dose:

- Eight (8) drops of FAITH™ Immune Fortifier.
- Add to these drops 8 drops of **ACTIVATOR**
- Continue with the protocols as for the previous doses

If no nausea was experienced continue with the protocols as follows:

Day 1 - Seventh dose:

- Nine (9) drops of FAITH™ Immune Fortifier.
- Add to these drops 9 drops of **ACTIVATOR**
- Continue with the protocols as for the previous doses

If no nausea was experienced continue with the protocols as follows:

Day 1 - Eighth dose:

- Ten (10) drops of FAITH™ Immune Fortifier.
- Add to these drops 10 drops of **ACTIVATOR**

If no nausea was experienced continue with the protocols as follows:

- Continue with the protocols as for the previous doses for Day 2.
- Continue increasing the drops daily with each dose until you reach a maximum of 10 drops per dose every hour or until nausea level is reached.
- Once nausea is experienced resume dosage at the previous level where there was no nausea experienced.
- Follow Aggressive Faith Formula Protocol until patient has recovered.
- Then resume treatment following the Maintenance Formula Protocols (see below)

THE MAINTENANCE FAITH™ FORMULA PROTOCOL

The Maintenance FAITH™ formula is as follows:

Follow the protocols for mixing the FAITH™ drops and do not exceed a total of three (3) drops ONCE daily for not more than ten (10) days per month.

Alternatively, administer three (3) drops of FAITH™ three (3) times a day at the onset of flu or if you experience any feelings of being unwell.

Drink this solution IMMEDIATELY it is mixed.

Recommended dosages for Children:

It is perfectly safe to have children and babies to take FAITH™.

Children should have fewer drops than adults.

For children 12 years and younger no more than TWO drops in total – starting with one drop, twice daily.

For children 13 years and older

- The formula is ONE drop per 10kg of weight not exceeding 3 drops per dose.
- Do not exceed this dose and not more than 3 times daily.

Larger doses.

Do not allow yourself to be fooled into doubling or tripling the doses. It does not mean that if 15 drops is good (per dose) that perhaps 45 drops per dose is three times as good. This is not necessarily the case.

DO NOT EVER BEGIN WITH MORE THAN 3 DROPS

Exception to the rule.

- In the event the person has a life-threatening disease only then could one start off with a large dose such as 15 drops.
- A large dose of drops often assists recovery in patients in the event of one having been poisoned, if one has had food poisoning or if one has been exposed to an infectious disease.

What time during the day is best to take FAITH™?

If you have recommended one dose per day of FAITH™, it is best to take just before going to bed as it works very fast and as a result, people often become sleepy after taking a dose of FAITH™. This simply indicates that the immune boosting process has begun. This sleepiness is a very good sign so allow them to take a nap if possible, for it is during sleep that the healing usually begins and they often feel much better when they awaken.

If your recommendation is to take it twice in a day, take one of the doses in the evening before going to bed. More aggressive regimes require multiple doses – ensure that the last dose is taken in the evening before going to sleep.

However, if you feel a cold coming on in the morning or in the afternoon do not wait until bedtime to take a dose. Take a dose anytime you feel like something is trying to come on.

Side Effects:

The immune system attacks unhealthy conditions using the FAITH™ drops and to date there has never been any evidence of an attack on healthy cells. However, if one supplies the immune system with excessive FAITH™ Drops, the immune system appears to attack unhealthy conditions vigorously but a nauseous reaction sometimes occurs, occasionally even vomiting and diarrhoea, this presents itself in 1:500 cases. When this does occur, one has simply taken too large a dose at first. Note that when you kill a lot of bacteria, viruses or parasites that have poison in them, that poison is likely to be dumped into the system and the end result is usually nausea and often even vomiting or loose stools. Simply reduce the number of drops.

One may also experience isolated pains related to the illness the person taking the drops has. This is as a result of the FAITH Drops assisting the immune system in attacking the pathogen / virus / bacteria etc. Do not stop drinking the drops at this point, rather continue, as the pain will disperse once the condition has been rectified by the body.

Please note: normally healthy people can take these drops without nausea, especially after a meal.

Stabilised Chlorine dioxide (SCD) is more powerful than any drug by far when utilised in the body to boost the immune system. The immune system selects and attacks many things that are not natural to the body and immediately destroys poisons that have been generated by disease. With the addition of the 12 herbal extracts (18 compounds), FAITH™ Drops has become an effective and powerful tool to ensure the removal of all of these toxins generated by disease as well as any toxic residue of the Chlorite used to generate SCD.

All known diseases and negative medical conditions:

We have not listed a great number of diseases which can be handled by the immune system in hours, days, or only a week or two, as there are so many. FAITH™ will probably help the immune system do its job no matter what the disease. For easy reference refer to the dosage table in Chapter 9 of this manual.

Diseases that are not caused by viruses, bacteria, and other germs may not be attacked directly by the FAITH™ but the benefit may come indirectly. Almost any disease generates various poisons which in turn cause the body to be sick.

When the immune system has SCD (Stabilised Chlorine Dioxide) which, of course, is furnished by FAITH™ it can often neutralize such poisons. Never assume that FAITH™ will not improve the immune system no matter how weak a condition the Immune system might be in or no matter what disease may be present itself in the body. It is not the disease that FAITH™ Drops is fighting – the drops are merely a tool to be utilized BY the body.

Take FAITH™ daily to keep the body clear of most poisons and disease-causing organisms.

Diseases that the immune system cannot help directly:

The immune system can use the FAITH™ to attack diseases caused by bacteria, viruses, parasites, yeast and moulds, but many diseases are caused by other medical conditions. Perhaps this is fact but since the first lot of research done, hundreds of people have reported almost every medical condition known as having been improved. There are no known diseases that do not respond to FAITH™ in some measure. Some diseases need to be handled by a proper diet and the immune system, when supplied with FAITH™ almost always produces some improvement in these diseases and their associated conditions.

Patients taking the drops and presenting with Lupus, Fibromyalgia, Diabetes, Depression and even ADD have reported improvements in their condition or in some cases complete recovery.

To ensure that the immune functions at optimum levels, we recommend that one makes use of the Maintenance Formula Protocols. A weakened immune system allows bacteria and viruses to infiltrate the weakened areas of the body and can often cause illnesses such as pneumonia etc.

With FAITH™ the immune system can fight back and often overcome the infections or other conditions that have been caused by the weakened condition. This can in turn help other diseases to clear up.

Remember, Stabilised Chlorine Dioxide is the most powerful killer of pathogens known to man. To ensure the safe removal of toxins, we have added 18 compounds (12 herbal extracts) to assist the body in the removal of any toxic residue.

FAITH™ can also be used Transdermally, anally and intravenously either with drip or direct injection. For these protocols contact the supplier directly, and / or refer to the dosage table in Chapter 9 of this manual.

Please note that in order to administer the drops intravenously, the person must be a licensed healthcare practitioner who has the permission of the MCC to administer the drops.

We at FAITH™ do not claim to heal anyone or anything. We only claim that our Immune Fortifier is exceptionally effective and provides the body with the necessary tools to do the job itself.

The immune boosting capabilities of the herbal extracts used in our formula are well documented.

Warnings:

1. Do not allow children to use FAITH™ unsupervised,
2. Do not allow FAITH™ to sit in direct sunlight even if it is in a colored bottle as it will create pressure in the container which could rupture and cause severe burns.
3. FAITH™ that has been sitting in direct sunlight can cause very painful burns.
4. Never allow full strength FAITH™ to remain on your skin for more than 10 seconds as it can cause mild burns.
5. Ensure that the product you use carries the official FAITH™ Drops Trademark to ensure that you do not cause additional damage to organs by drinking generic copies of our product.
6. Please ensure that should you venture upon a "generic" version of our product that you request from the supplier copies of the products NAPPI Code, the registration documents with the Medicine Control Council as well as the Consumer Council registration Documents. If you are still not sure that it is FAITH™ Drops that your have been sold, check for the call centre number on the labels and boxes as well as batch numbers and expiry dates. Generic copies of our product are currently being manufactured at home WITHOUT the essential 12 herbal extracts that protect the body from toxin. The supplier of this generic product will not be able to show you documentation of the pharmaceutical company that is authorized to manufacture their product under licence. The manufacturing company should be registered with the Pharmaceutical Council of South Africa. If after all this you are unsure, FAITH™ has a very sophisticated consumer and agent registration database – we can tell you with ONE phone call if the person selling you this product is a registered agent or NOT. If the agent is registered it means they have undergone training in the use and application of the FAITH™ Drops and have been accredited by the manufacturer of the product.

Using FAITH™ on the Outside of the Body:

Please read this carefully.

It is possible to cause problems on the outside of the body when using FAITH™ in too strong a solution.

Once you have added the activator to FAITH™ Drops it can no longer cause a burn, but it can still cause a problem when used full strength.

- When adding six to ten drops in a glass with 6 – 10 drops activator, with the exception of insect bites covering small areas of the body, always add at least 125ml (125cc) of water before putting the diluted solution on the skin or in the hair. For specific dosages refer to Chapter 9.
- Never add the 10 drops mixture to any kind of body lotion.
- When the FAITH™ solution is too strong on the outside of the body, it can weaken the cells and make them more susceptible to bacteria and infection.

When the problem doesn't clear up right away:

SCD is an exceptionally strong oxidizer and its purpose in the body is to oxidize any unhealthy deposits in critical areas that are making you ill. Poisons such as heavy metal deposits or bacteria growing somewhere or any other unhealthy conditions will need to be worked on continuously by the immune system. However, the few drops that one takes daily will result in only a few milligrams of SCD entering the system and it is this small quantity of SCD that has to aid the body to remove pathogens. This could be a long slow process or one could see or experience a difference in a very short space of time. Each person is different.

If you are experiencing nausea when you try to increase your dosage level this should indicate to you that the immune system is coping with toxins in the body. As advised already, reduce the number of drops to just below nausea level. Remain at that level until you are able to increase the drops – slowly. Some people are able to immediately begin with 3 drops and increase daily until they are taking 15 drops per dose without any problem whatsoever. Others might find it difficult to begin with even one drop. As explained each person is different and perseverance is the key to success.

Diabetes, ALS and many other diseases

The Coxsackie's B virus, according to much of the present medical literature, causes up to 1/3 of the heart attacks. It causes the heart to become inflamed. The heart doesn't work very well when inflamed. It is also reported that this virus causes the pancreas to become inflamed thus causing type 1 and type 2 diabetes.

In the case of ALS (Lou Gehrig disease) there is medical literature indicating the echo virus - 6 and - 7 is most likely the cause. If that is so, then there is a chance that FAITH™ can help cases of ALS.

The idea is to keep the body as saturated as possible with the SCD.

There is no guarantee, but up until now there has never been even a slight glimmer of hope.

Curing diseases with drugs?

Can it be done? No. The only thing that ever heals the body is the various mechanisms of the body. Healing is controlled by the body. NO drug AND no nutrient can heal the body. There are things that the body can use to aid in the healing process, but in the final analysis, the body always does the healing.

FAITH™ cannot heal or cure the body of anything. It may or may not kill pathogens in the body without the body's control. However, once the germs, pathogens or poisons are destroyed, healing can take place at a much faster rate. This is why healing in so many parts of the body seems to happen so much quicker. It is not that FAITH™ has healed anything, it cannot - but it can clear the way for healing to take place.

We have every confidence in our product, however, what we don't have confidence in is mans impatience with himself and the possible factor of human error.

In cases of chronic diseases such as cancer, hepatitis, heart conditions and diabetes, we recommend that you consult with your healthcare practitioner before beginning the drops and that you continue with your regular check-ups and medication.

Our Head Office will be able to refer you to a Doctor that is accredited in the use and application of FAITH™ Drops. Please feel free to call them at any time.

- Always reduce the number of drops if you feel nausea, but continue to increase the number of drops, keeping below the nausea level as much as you can.
- **Do not** remain at high dosages unless you really see an improvement in the condition you are trying to handle.
- When the condition is overcome, be sure to drop back to make use of the Maintenance Formula Protocols.

Antidote for too much FAITH™:

- For someone who has taken too large a dose of FAITH™ and is then nauseous, it is possible to drink a glass of cold water to which has been added 1000 MG of vitamin C.
- Bicarbonate of soda also as such an antidote. Use a level teaspoon full of Bicarbonate of Soda or "Enos" in a glass of water.
- If you use the vitamin C do not use the Bicarbonate of Soda or "Enos".
- Use either one but not both.

Overdose of FAITH™:

- Anyone who has consumed more than 2.5ml (2.5cc) of FAITH™ directly from the bottle – inactivated and undiluted one should immediately drink as much water, as much as possible.
- It is possible to drink as much as several tablespoons of FAITH™ straight from the bottle, but this would make one very sick, and is NOT the way the drops are intended to be utilised.
- It is best to add ½ teaspoon full of Bicarbonate of Soda to each glass of water. Be sure to see a doctor or emergency poison clinic as soon as possible.

Training the individual:

You as the healthcare practitioner have read these notes and hopefully you now know how to use the FAITH™ drops BUT if you have recommended that your patient take the FAITH™ drops, YOU must train them to use it properly. Do not expect that you can rely only on the directions on the insert. People must be trained to use the drops by actually holding a training session.

Have the individual drop two drops of FAITH™ Drops into an empty glass, add an equal number of drops of ACTIVATOR. Wait the prescribed 3 minutes then add a MIXER, such as water.

Every mistake in mixing the drops has been made to date – mostly because people either don't listen, don't read or don't understand. If you sit with them whilst they mix their first dose and they make their own notes – they cannot make a mistake.

Making mistakes in mixing the formula only hinders their recovery. Train them properly – today.

Please stress to the user that they should ALWAYS read the bottle labels and leaflet/inserts and should err on the side of caution if they are still unsure of how to proceed. Make sure that the user is able to understand the importance of this.

ALWAYS READ LABELS AND INSERTS. MAKE SURE YOU UNDERSTAND HOW TO PROCEED.

REFER ANY QUERIES TO YOUR HEALTHCARE PRACTITIONER OR EMAIL US ON: damaansax@vodamail.co.za

Observations:

The following information has been discussed with medical doctors who have agreed with me, but some had their own opinion as to what FAITH™ actually does in the body. These notes are from ten years of observing people that are recovering whilst taking FAITH™ drops.

What appears to be happening in the body when FAITH™ is ingested after using the various activators listed are:

- 1). FAITH™ Drops destroys anaerobic microorganisms including viruses, bacteria, molds, yeast, and parasites. These are the disease causing organisms. There are many reasons to believe that this is what happens. This generally happens between 4 hours and 4 weeks, but in less than one week in most cases. This includes all forms of infection and other microorganisms, as well as blood diseases such as leukemia.
- 2). FAITH™ Drops oxidizes the heavy metals within the body. We believe this because there have been a number of tests where the hair roots were checked before and after taking FAITH™. Afterwards, the heavy metals were gone. That includes mercury and lead and several other metals. Unfortunately, hair root testing is not totally accurate, but it is a good indication. Oxidation of the heavy metals is not the same process as chelating, but the results are the same. When a heavy metal is oxidized it is neutralized and is simply washed out of the body.
- 3). FAITH™ Drops is carried throughout the body where it neutralizes foreign matter that is generally poisonous to the body. When oxidized, these poisons are neutralized and merely wash out of the body. Almost all poisonous material will be acidic in nature, thus allowing FAITH™ to recognize it. These poisons will all be attached at various locations and they will be hindering the function of the body. As they are oxidized they will no longer stay attached and will *be* washed out of the body.
- 4). Most forms of poisons from snakes and other venomous animals are oxidized when adequate amounts of FAITH™ Drops is taken. Most food poisons found in restaurants or in your refrigerator are oxidized, which is why one should take a number of doses when feeling the effects of food poisoning.
- 5). The poison generated in burns, particularly type 3 burns when covered immediately with FAITH™, or even within hours, is neutralized by the FAITH™ Drops. The FAITH™ Drops solution should not be left on more than 1 minute and it should be rinsed off with sterile water.

Using FAITH Intravenously

For making use of FAITH™ intravenously, kindly contact Dr. Michael McDonald via email.

Anyone attempting to perform intravenous applications without first being a registered medical practitioner and without MCC permission does so at their own risk.

Dr. Michael McDonald

Email: damaansax@vodamail.co.za

Cell: +27823399799

Overview of the Use of FAITH and Main Points

1. A maintenance dose of FAITH™ is three drops once a day for ten days in one month.
2. A full dose to overcome diseases is maximum 10 drops per dose. This is for Chronic Diseases or for life threatening disorders.
3. For life-threatening disease a full dose would be administered every hour each day until the disease is gone. Other protocols may apply for certain diseases. See table in Chapter 9.
4. One drop of FAITH™ contains nine milligrams of sodium chlorite.
5. When six drops of FAITH are mixed with six drops of **ACTIVATOR** in a dry clean glass, one milligram of chlorine dioxide will be generated in three minutes. The solution will not become much stronger than one milligram as the chlorine dioxide near the surface of the liquid goes into the air. You can make it stronger by putting a lid on the solution preventing much of the chlorine from escaping.
6. When 125ml (125cc) of apple juice, grape juice, or pineapple juice **MIXER** is added to the mixture and the drink is consumed, one milligram of chlorine dioxide will be generated each hour for approximately 12 hours within the human body.
7. Chlorine dioxide, generated by FAITH™ Drops, is the most powerful killer known to man of microorganisms, such as, viruses, bacteria, parasite, molds and yeasts.
8. Chlorine dioxide has been known and used for over 100 years as a sterilizer for hundreds of uses in hospitals and the food industry.
9. Do not allow FAITH™ to sit in direct sunlight. It creates high pressure and has been known to blow the lid off. The liquid after contacting sunlight becomes very strong and will create burns if not washed off immediately. In the case of contact with eyes, flush with water for several minutes.
10. Any dose of FAITH™ requires an equal number of drops of **ACTIVATOR** in the form of citric acid **with the exception of intravenous use**. Without citric acid there is very little benefit.
11. Do not use **ACTIVATOR** (citric acid) when used intravenously. The blood will accomplish what the **ACTIVATOR** would do otherwise, but it does it more slowly.
12. Only registered Medical Practitioners may administer the drops intravenously and only with the express permission of the MCC.
13. Sodium chlorite is used throughout the world to generate chlorine dioxide for water purification.
14. Sodium chlorite was first used for treatment of humans in 1926 in Germany by Dr. William F. Koch, MD, Ph.D. It has been use in the U. S. since 1930.
15. Since 1926 it has been considered by those who use sodium chlorite that it is Stabilized Oxygen and that it furnishes oxygen to various parts of the body. That is a mistaken idea. No useable oxygen is available from sodium chlorite; however Chlorine dioxide is also an oxidizer and is much more powerful than oxygen.
16. Vitamin C that is used in juices as a preservative will completely prevent the FAITH™ from working. **DO NOT USE JUICES THAT HAVE VITAMIN C ADDED TO THEM. READ THE INGREDIENTS ON THE CONTAINER FIRST.**
17. Some pure apple juices do not have added vitamin C.

18. Orange juice will prevent FAITH™ from working. It prevents the chlorine dioxide from being generated.
19. Any of the following **MIXERS** - Apple juice, cranberry juice, grape juice, and pineapple juice are okay to add to the FAITH™ formula, after the three minute wait - if the juices have not had vitamin C added.
20. Shelf life, if the bottle has not set in the direct sunlight, is two to four years if the bottle is dark glass or plastic. If the bottle is in the direct sunlight it will last only about an hour even when if it is in dark glass or plastic.
21. Full strength FAITH™ Drops is alkaline at a pH of 13. It can cause a slight burn if not washed off of the skin in less than a minute. Flush with water. When activator is added to 6 or 12 or even 15 drops the pH is then 4.5. When apple juice is added after three minutes the pH matches the apple juice exactly, which is approximately 4.8. The other juices mentioned in this book are similar. Use only apple, grape, pineapple or cranberry juices that do not have vitamin C added.
22. FAITH™ Drops can be used to purify water. Use 8 drops per gallon when in the jungle or in the woods. Wait 8 hours after adding the drops. The water is then safe. Or use 6 drops per gallon in a foreign country to make the water safe.

ADDITIONAL RESEARCH

The Overnight Cure for Cancer (OCC)

by R. Webster Kehr of the Independent Cancer Research Foundation, Inc.

Disclaimer

This treatment has not been evaluated by the FDA. The term "cure," as used in this article, is the alternative medicine definition of "cure," meaning the vast majority of the cancer cells are killed or reverted into normal cells.

General Warnings

This cancer treatment is new. It is experimental, though it has been proven to be totally safe even for very advanced cancer patients. In other words, its safety is NOT experimental, only its effectiveness is experimental. This treatment is based heavily on a great deal of scientific evidence combined with solid cancer theory.

Be aware that chlorine dioxide is toxic in high doses. The doses chosen for this treatment have been chosen to be well within safe, non-toxic levels. However, do not assume significantly higher doses of chlorine dioxide will be safe. Stick within the safe doses of this treatment. Several people have already taken this treatment and no one has had the slightest complaint about the doses.

Please note that this treatment is spread out over 12 hours.

This is also part of the safety of this treatment. Do not take the total daily doses in less time than the treatment prescribes!!

Note further that DMSO is far, far less toxic than chlorine dioxide, thus the doses of DMSO are simply not a factor. Earlier versions of this treatment used 25 tablespoons of DMSO without a single complaint. This treatment uses less than ½ teaspoon of DMSO during the entire treatment.

Because this is a new treatment, this article changes from time to time. It is important to read this article in its entirety just before starting the treatment (i.e. after you have obtained the necessary materials). This article is still changing as feedback is received from cancer patients who have taken the treatment.

The OCC is Not Ready for Advanced Cancer Patients

It will take several months before the OCC can be fine-tuned to be effective for advanced cancer patients. Thus, the Overnight Cure for Cancer should NOT be used by advanced cancer patients. This is not because the OCC is dangerous; it is because advanced cancer patients should not take a week off for any cancer treatment which has not been proven to be effective for advanced cancer patients.

In other words, advanced cancer patients should ONLY use alternative cancer treatments which have been proven to be effective for advanced cancer patients, such as the Collect-Budwig Protocol or the Bill Henderson Protocol or the Cesium Chloride Protocol.

See the Cancer Tutor website for a discussion of the proven alternative cancer treatments.

Children under Twelve Should NOT use This Treatment except under the supervision of a Medical Professional.

Do NOT Use This Treatment whilst using Prescription Drugs.

This alternative cancer treatment should NOT be combined or used with ANY prescription drugs. The reason is that this treatment may greatly enhance the effectiveness of prescription drugs. For example, if a prescription drug binds to DMSO, which is part of this treatment, the DMSO may drag the prescription drug into the cancer cells, thus killing the cancer cells. This treatment is designed to transform cancer cells back into normal cells. Killing cancer cells may yield undesirable inflammation and swelling and/or create excess debris from dead cancer cells.

Do NOT Use This Treatment with some Alternative Cancer Treatments

There are two kinds of alternative cancer treatments or products which should NOT be used with this treatment.

1. This cancer treatment should NOT be combined with alternative cancer treatments which kill cancer cells. Should a product which kills cancer cells bind to the DMSO, the DMSO may allow it to kill far more cancer cells than it normally would. This could lead to excess debris from dead cancer cells. In other words, do NOT combine this treatment with any alternative cancer treatment or product which kills cancer cells!!

This treatment is so potent; if cancer cells were killed it could create a dangerous situation for the patient!! Use either DMSO combined with Faith or select the alternative treatment of your choice BUT DO NOT combine the two treatments.

2. This treatment should NOT be used with alternative cancer treatment products which contain Vitamin C, Vitamin E, selenium or any other antioxidant or immune builders. These products will neutralize the effectiveness of this treatment. More will be said about these products below.

Warning For Women Who Are, or Might Be, Pregnant

Women who are pregnant, might be pregnant, might become pregnant, or are nursing, should NOT take this treatment. The affect on an unborn fetus could be fatal to the fetus due to the high doses of chlorine dioxide in this treatment combined with the extremely low weight of the fetus!! In addition, fetus have many undifferentiated cells and this treatment will TARGET cancer cells, which are also undifferentiated!! Thus, this treatment may inadvertently target undifferentiated fetal cells!!!!

TAKE THIS WARNING VERY, VERY SERIOUSLY!!!

ALLERGIES TO DMSO

I have never seen a case of an allergy to DMSO, but apparently in rare cases someone is allergic to DMSO. When you get your bottle of DMSO put one drop on your skin and spread it out and see if you have an allergic reaction. If not, an hour later put 10 drops on your skin and spread it out. If you do not have a reaction, go ahead with the treatment.

What Causes Cancer?

Most people believe that it is DNA damage that causes cancer. While in rare situations, DNA can have a negative affect on a person's immune system. DNA normally has absolutely nothing to do with the development of cancer.

The "theory" that DNA causes cancer is driven by a desire to convince the general public that a cure for cancer is 100 years away. This deception is to convince the general public that cancer can only be treated as a highly profitable chronic disease. The cancer treatments and propaganda which are chosen by the FDA and pharmaceutical industry (and world-wide for that matter, I know of no country on earth with honest politicians) are chosen on the basis of profitability, not effectiveness. There is no money in curing cancer, but there are hundreds of billions of dollars in pretending cancer is a chronic disease and there is no cure for cancer. All of this is nonsense driven by greed.

The fact is that cancer is caused by a special type of microbe which gets inside of normal cells and turns the cells cancerous.

Dr. Royal Rife did an enormous amount of research into the relationship between microbes and cancer in the 1930s. He would inject mice with a virus and in 100% of the time the mice would get cancer.

Dr. Rife proposed a cure for cancer which did nothing but kill these viruses. His cure was successful. However, note that his cure had no intention of killing cancer cells; its only goal was to kill microbes which were inside of the cancer cells. Once the microbes were dead the cancer cells were able to revert back into normal, differentiated cells.

Many other cancer researchers, starting over 100 years ago in the 1800s, have isolated the cause of cancer to be microbes, though they did not understand the mechanism inside the cell which caused a microbe to make a cell cancerous.

Now, the entire mechanism inside the cell which allows microbes to cause cancer is understood. We now know that a microbe which is able to get inside of a normal cell blocks glucose from being used to create pyruvate, which in turn blocks the Citric Acid Cycle and in turn the Electron Transport Chain, both in the mitochondria. Blocking these two chemical chains cause the number of ATP molecules in a cancer cell to plummet!!

A detailed discussion of how microbes cause cancer can be found in the Cancer Theory article. In essence, the article on cancer theory is also the theory behind this treatment. The cancer theory article also discusses the four different categories of treatments which can cure cancer. See this article:

Cancer Theory.

You may have noted in the article just linked to that as long as microbe(s) are inside cancer cells, the cell is unable to revert into a normal cell.

While many people have tried to replicate Dr. Rife's electromedicine cancer treatment (which was destroyed by the FDA and AMA), this treatment, the Overnight Cure For Cancer, is the first non-electromedicine cancer treatment ever designed to specifically kill the microbes inside the cancer cells as a cure for cancer.

If you kill the microbe(s) inside the cancer cells the cancer cells WILL NOT DIE!! The cancer cells will actually be able, within a few days, to restore their Krebs Cycle and Electron Transport Chain and become normal, differentiated cells again. Thus, there is zero debris from dead cancer cells or broken-apart DNA. That is why this treatment can be so effective so fast.

This treatment is not only based on solid cancer theory, it has behind it a great deal of scientific evidence.

So how is it possible to kill a microbe which is comfortably living inside a cancer cell?

First, you must get a chemical which is known to kill microbes inside the cancer cell. Chlorine dioxide is such a molecule.

So how do you get chlorine dioxide inside the cancer cells?

You use DMSO. DMSO is officially: Dimethyl Sulfoxide. DMSO is a byproduct of the wood industry and is totally natural and will bind to many different kinds of molecules and drag them through the skin into the bloodstream (i.e. transdermally).

The details of the necessary sequence for the Overnight Cure For Cancer (OCC) to work is as follows:

1) You find a molecule known to kill microbes on contact chemically, not mechanically. Chlorine dioxide is such a substance. In fact, chlorine dioxide is the substance used in the OCC to kill the microbes inside the cancer cells. Because chlorine dioxide normally stays in the bloodstream, it generally only kills microbes in the bloodstream.

2) DMSO must be able to bind to chlorine dioxide so that the chlorine dioxide will get into the bloodstream transdermally. Taking chlorine dioxide orally can be a very slow process; taking weeks or months to build up to therapeutic doses. DMSO has been studied and used since the early 1960s to bind to molecules and drag them through the skin into the bloodstream.

3) Once inside the body, the DMSO, with chlorine dioxide in tow, must target cancer cells. This is a very well-known ability of DMSO. DMSO has been combined with chemotherapy, and many other substances, to get them inside of cancer cells. The DMSO/chemotherapy cancer treatment, which was designed to kill cancer cells by using DMSO to get chemotherapy inside the cancer cells, was a superb cancer treatment, but it was shut down by the FDA in Georgia.

4) Once at the site of a cancer cell, the DMSO and chlorine dioxide must get inside the cancer cell.

5) Once inside the cancer cell the chlorine dioxide must be able to kill the microbe(s) inside the cancer cells. Once the microbe(s) are killed inside the cancer cell, the cell will be able to restore its Citric Acid Cycle and Electron Transport Chain (ETC). In about a week the cancer cell will be a normal cell. The cell will revert from an undifferentiated cell to a differentiated cell.

Most of the above items are known to be true. For example, it is KNOWN that DMSO and chlorine dioxide will bind together. And it is KNOWN that DMSO will carry chlorine dioxide through the skin.

It is also known that chlorine dioxide can kill microbes while bound to DMSO. For example, I sprayed some chlorine dioxide, bound to DMSO, on some mould. Within a minute the mould was dead. Chlorine dioxide bound to DMSO has also been used to cure toenail fungus, thus we know DMSO and chlorine dioxide will penetrate skin, and while bound together, chlorine dioxide can kill microbes.

In fact, the only issue which has not been proven is that DMSO and chlorine dioxide, bound together, can get inside a cancer cell. However, knowing how cell membrane ports are built using proteins, and knowing the size of the DMSO / chlorine dioxide molecule, it seems obvious that they will get inside the cancer cells.

Furthermore, as just mentioned, DMSO is known to pull some forms of chemotherapy inside of cancer cells. Thus, it is highly likely it will pull chlorine dioxide through the cancer cell ports.

However, a new issue which has recently come to light is whether the diet of the person may neutralize, to some degree, the effectiveness of the chlorine dioxide itself.

This treatment is very different than the normal way that DMSO and chlorine dioxide are used. The reason for the intensity of this treatment is that it is necessary to kill EVERY microbe inside of EVERY cancer cell in a short amount of time. If you don't accomplish that, the surviving microbes will continue to breed inside the cancer cells and the cancer will again start spreading. So it is a safe, but intense, treatment.

Why Is This Called the: Overnight Cure for Cancer?

The vast majority of alternative cancer treatments kill cancer cells. Cancer treatments which kill cancer cells can only kill so many cancer cells per day or else the debris from dead cancer cells can overwhelm the liver.

While chemotherapy must be "paced" because of all the damage done to non-cancerous cells; many alternative cancer treatments must be "paced" because they target cancer cells and kill so many of them that the liver is burdened with debris from dead cancer cells.

Thus, most alternative cancer treatments must be "paced," meaning doses of substances must be limited to the ability of the liver to process the debris from dead cancer cells.

Some of the treatments that kill cancer cells also cause swelling and inflammation; even tumors can swell during the treatment. The reason for this is that if cancer cells are killed slowly, the immune system will recognize the cells as being "sick" and will attack them. The inflammation, swelling and possibly congestion (for lung cancer) can also cause alternative cancer treatments to need to be paced.

For example, the normal dosage for cesium chloride and DMSO must be cut in half for brain cancer and certain other cancer conditions.

However, when you kill the microbe(s) inside a cancer cell, but do NOT kill the cancer cell itself, the immune system is clueless as to what is going on, thus there is NO inflammation or swelling.

But equally important, when you revert a cancer cell into a normal cell there is no debris from dead cancer cells or broken apart DNA. The cell stays intact and the cell deals with the debris from the dead microbes "in house," so to speak. The debris which goes outside the cell is minimal because the cancer cell itself is not killed.

Thus, in theory, a treatment which reverts cancer cells into normal cells could cure cancer within 24 hours!! But the only way to do that would be to kill the microbes inside the cancer cells without killing any cancer cells! That is exactly why the Rife Machine could safely cure cancer within minutes, without any danger to the patient.

Excluding electromedicine, there are more than a dozen natural substances which are known to be able to revert cancer cells into normal cells. However, none of them are practical as a cancer treatment for a variety of reasons.

But this treatment is practical, and that is why this treatment is called the "Overnight Cure For Cancer." When this treatment is perfected, it literally will be able to cure cancer within 24 hours, without any debris from dead cancer cells, debris from broken DNA or swelling or inflammation from the immune system attacking cancer cells which are slowly dying.

The OCC is also safe for the liver, unless the liver has already been damaged by debris from dead cancer cells or chemotherapy.

The importance of perfecting this treatment cannot be overestimated. It overcomes all the problems of other alternative cancer treatments.

A CRITICAL, CRITICAL CONCEPT!!!

Curing cancer, especially advanced cancer, is in two parts. They are both equally important!!

The two parts are the "cancer cells" and the "non-cancerous cells." The importance of the cancer cells is obvious. So why are the non-cancerous cells just as important as the cancer cells when treating cancer?

Many cancer patients, perhaps most cancer patients, die because of the damage done to their non-cancerous cells. Cancer cells steal vital nutrients from non-cancer cells, that is one reason cancer patients become so weak. In addition, chemotherapy and radiation do incredible damage to non-cancerous cells. Thus, in advanced cancer, the non-cancerous cells are very, very weak and very, very sick.

The Overnight Cure For Cancer will ONLY deal with cancer cells, it WILL NOT help the non-cancerous cells.

For example. Suppose there was a "magic bullet" (which hopefully the OCC will become) which safely rid the body of all cancer cells within one day. Would the cancer patient be guaranteed to survive their cancer? The answer is 'no', because the non-cancerous cells are the cause of death in many cancer patients, if not most cancer patients. Thus, getting rid of the cancer cells is only half of the battle with cancer.

So here is the key question. Should a cancer patient deal with the cancer cells first or the non-cancer cells first?

This question will be answered with a parable:

The Parable of the Thugs and the Preachers

Suppose there were 100 thugs who were locked in a building with 1,000 preachers. Suppose at every opportunity the thugs beat up the preachers. The preachers were getting weaker and weaker because of getting beat up daily by the thugs.

You are assigned to go in and help nourish the preachers back to health. What is the first thing you should do?

The first thing you should do is get rid of the thugs!! If you don't get rid of the thugs you will not be able to nourish the preachers because they will continue to get beat up as you are trying to help them.

In exactly a similar way, it is critical to get rid of the cancer cells as quickly as possible!!

However, you cannot kill the cancer cells quickly or it will create so much debris you may kill the cancer patient.

But the Overnight Cure for Cancer does not kill cancer cells, not a single one. The OCC is designed to revert cancer cells into normal cells very, very quickly.

- **This treatment should be used three times, until more is known about its effectiveness.**
- **There should be a two day period BETWEEN each of the three times this treatment is taken.**
- **The pre-OCC, which is taken the night before the actual OCC, can be counted as one of the two rest days.**

Thus, the schedule will look like this:

Day 1) Pre-OCC Day (4 "doses" taken the night before the OCC). This day will be discussed below.

Day 2) The complete OCC

Day 3) Rest and eat healthy foods to nourish the non-cancerous calls.

Day 4) Rest and eat healthy foods to nourish the non-cancerous calls, but also take the pre-OCC

Day 5) The complete OCC

Day 6) Rest and eat healthy foods to nourish the non-cancerous calls

Day 7) Rest and eat healthy foods to nourish the non-cancerous calls, but also take the pre-OCC

Day 8) The complete OCC

After this, you should go on other alternative cancer treatments.

During the time you are buying the items in the OCC you should also be ordering the items in the Collect-Budwig Protocol, the Bill Henderson Protocol or whatever alternative cancer treatment you choose.

Taking Chlorine Dioxide: Oral or Transdermal?

As mentioned above, Vitamin C, other antioxidants and immune builders, should not be taken during the same time periods as MMS/Chlorine Dioxide. This is good advice.

However, most websites recommend chlorine dioxide be taken orally. When taken orally, the treatment can cause diarrhea, nausea, vomiting, etc. The side-effects can sometimes be so severe it can take many weeks for a patient to build up to a therapeutic dose. These side-effects only happen if you take this treatment orally. It is the stomach which is causing these side-effects. If you take this treatment transdermally, meaning through the skin; the stomach, and its side-effects, can be totally avoided.

Due to the volume of chlorine dioxide which will be taken in a short amount of time, taking chlorine dioxide orally is absolutely not an option for this treatment.

In addition, since the DMSO is so very critical to this treatment, the DMSO will also cause some stomach upset if taken orally. Thus, there is a second reason to take the treatment transdermally.

The most important advantage of taking it transdermally is that with transdermal applications, higher doses can be taken (actually small doses are taken several times), and the high daily doses can be achieved much more quickly.

It is rare when more than 30 drops of chlorine dioxide are taken in a day. In this protocol, 130 drops, depending on the weight of the patient, are taken. The drops are spread out evenly over 12 hours (it is critical to spread out the drops). Thus, this is a different kind of treatment.

The Substances Needed For the OCC

Here is a checklist of things you need for this treatment.

1. FAITH DROPS - At least 4 bottles
2. ACTIVATOR
3. 500ml DMSO – diluted with 125ml of distilled WATER – then decant into storage bottles.
4. MSM crystals to be made into "MSM Water" (see below).

DO NOT use a DMSO gel or DMSO cream. These will not bind to chlorine dioxide!!!

The purpose of the MSM is to prevent the DMSO from breaking down inside the body. It is a very important part of the treatment!! MSM will penetrate the skin, just like DMSO, but it will NOT bind to other molecules or carry them through the skin.

In some cases a person can obtain all the substances needed for the OCC except for the MSM. The MSM Water is highly recommended for this treatment, but it is optional. In other words, don't let the lack of MSM stop the rest of this treatment.

Making "MSM Water"

Here is how you make MSM water:

1. Take 4 Litres of DISTILLED water
2. ADD 1 ½ CUPS (375ml) of MSM granules to the container of distilled water.
3. It will take about half-an-hour for the MSM to totally dissolve in the water.
4. Shake the container every few minutes until it is totally dissolved at the bottom of the jug. This is the "MSM Water."

How to Make "ONE DOSE" of Chlorine Dioxide and DMSO

AT ALL TIMES DURING THE MAKING OF YOUR DOSE OF CHLORINE DIOXIDE WHEN EVER POSSIBLE, USE A PURE SILVER TEASPOON. IF ONE IS NOT AVAILABLE, USE A STAINLESS STEEL TEASPOON BUT NEVER A PLASTIC TEASPOON.

How many times you take the dose of chlorine dioxide and DMSO will be discussed later. But first, we need to understand how to make "One Dose."

READ THIS SECTION SEVERAL TIMES to make sure you aren't missing anything. Especially read it **AFTER** making one or two doses!! You would also be smart to have a second person look at the instructions to make sure you both agree on how to make "One Dose."

Step 1: Put exactly 20 FAITH drops into a small glass bowl.

Step 2: Add 20 drops of activator to the FAITH in the small glass bowl.

Step 3: Your **FIRST** Wait of 3 Minutes.

AFTER MIXING the FAITH drops with the activator, stir the mixture using as already instructed a pure silver teaspoon if possible, but not plastic, then let it sit for 3 minutes. Stir the mixture every 30 seconds. This three minute wait creates the chlorine dioxide.

Step 4: Measuring the DMSO

AFTER the 3 minutes needed to make chlorine dioxide, add the liquid DMSO to the mixture.

In the same small bowl add (¾ of a **teaspoon**).**75cc** of DMSO. In other words, this is added to the chlorine dioxide you made above regardless of your weight.

Step 5: Your **SECOND** Wait of 3 Minutes

After **ADDING** the DMSO to the chlorine dioxide you need to mix the mixture, then wait an additional 3 minutes, stirring the mixture every 30 seconds.

This is the **SECOND TIME** you have waited 3 minutes. This time you are waiting for the DMSO to bind to the chlorine dioxide.

After this second wait of 3 minutes you have a mixture which is defined to be "One Dose" of Chlorine Dioxide / DMSO.

Clarifying What "One Dose" Means

The steps above can be summarized thusly:

- 1) Put 20 drops of FAITH in a small glass bowl,
- 2) Add 1 teaspoon of citric acid.
- 3) WAIT 3 minutes (stir at first and every 30 seconds) to make the chlorine dioxide,
- 4) Add $\frac{3}{4}$ of a **TEASPOON** of DMSO
- 5) WAIT an additional 3 minutes (stir at first and every 30 seconds) for the DMSO to bind to the chlorine dioxide.

You Have Now Made "One Dose"!!

Applying the DMSO dose Transdermally

- When the second three minute wait is completed, you can rub the entire mixture onto your skin.
- Spread the mixture so that it is thin on your skin. This way it will penetrate faster and will create less of a skin rash.
- It is very important to rotate where you put the mixture on the skin.
- During the "prep" day you will take "One Dose" 4 times, where each dose is separated by 1 hour.
- The first time you may want to put it on your arms.
- The second time you may want to rub it on your leg thighs.
- The third time you may want to rub it on your calf muscles.
- The next time you use it you will rotate back to your arms etc.
- The person's back can also be used.

By rotating where you put the mixture your skin has 3 hours to completely recover from the DMSO pulling the chlorine dioxide through the skin.

Having said all of these things, an even higher priority would be to put the mixture as close to the cancer as possible. As much as your skin can tolerate it, put as much of the mixture on the skin above where the cancer is located.

There are ways to protect your skin, no matter where you put the mixture. Ten minutes after putting the "One Dose" on the appropriate place on your body (if you can wait that long), you can put MSM Water on that location of the skin if there is any rash developing.

Thus, you have two different ways to protect your skin during the treatment; first, by spreading the mixture as thin and wide as possible, and second, by using MSM water ten minutes after spreading the mixture on the skin. But as mentioned before, put the mixture as close to the cancer as possible as much as you can.

It should be mentioned that latex gloves, rubber gloves or any other kind of gloves should NOT be used to spread any mixture containing DMSO. The DMSO can pull the materials in these gloves through the skin. Always use bare hands to spread the mixture.

Phase One of the OCC - the "Pre-OCC" - the Night Before the Main OCC

Note: If at any time during the "Pre-OCC Day" or the "OCC Day," if you do not feel well, terminate the treatment immediately. This treatment is not toxic, but there may be individuals who have a reaction to the DMSO or chlorine dioxide.

For most situations, the chlorine dioxide / DMSO protocol will be in two phases:

Phase One:

- This is the day before the actual Overnight Cure For Cancer (OCC).
- On this day you will take four doses of chlorine dioxide and DMSO, separating each dose by one hour.

You should take these four doses after dinner. Thus, you might take them at:

- 6:00 PM (1800)
- 7:00 PM (1900)
- 8:00 PM (2000)
- 9:00 PM (2100)

Also, at the one-half hour mark, between the one hour marks, you need to spread one **tablespoon** of MSM Water on the skin, especially where you had put the DMSO. This is in ADDITION to any MSM Water you used to prevent skin rashes.

Thus, your Pre-OCC schedule might be like this:

- 6:00 PM (1800) - One Dose of the Chlorine Dioxide / DMSO Mixture
- 6:30 PM (1830) - MSM Water ONLY
- 7:00 PM (1900) - One Dose of the Chlorine Dioxide / DMSO Mixture
- 7:30 PM (1930) - MSM Water ONLY
- 8:00 PM (2000) - One Dose of the Chlorine Dioxide / DMSO Mixture
- 8:30 PM (2030) - MSM Water ONLY
- 9:00 PM (2100) - One Dose of the Chlorine Dioxide / DMSO Mixture

This pre-OCC is critical!!! The purpose of Phase One is to kill the microbes in the bloodstream. If you do not do this, some of the chlorine dioxide / DMSO in Phase Two, the actual OCC, will be diverted to kill microbes in the bloodstream. This is not the goal of Phase Two. The goal of Phase Two is to kill the microbes INSIDE the cancer cells to revert the cancer cells back into normal cells.

Phase Two of the OCC - The Main OCC

- The actual OCC (i.e. Phase Two) consists of 13 doses of chlorine dioxide and DMSO.
- If you cannot manage the 13 doses, do the best you can.
- As before, they should be taken one hour apart.
- As before, at the half-hour mark between CD/DMSO doses, you should put MSM Water on your skin.

Let us assume you start at 8:00 A.M. Your schedule would look like this:

DOSE NUMBER	TIME	WHAT TO TAKE
1	08:00	One dose
1	08:30	MSM water ONLY
2	09:00	One dose
2	09:30	MSM water ONLY
3	10:00	One dose
3	10:30	MSM water ONLY
4	11:00	One dose
4	11:30	MSM water ONLY
5	12:00	One dose
5	12:30	MSM water ONLY
6	13:00	One dose
6	13:30	MSM water ONLY
7	14:00	One dose
7	14:30	MSM water ONLY

DOSE NUMBER	TIME	WHAT TO TAKE
8	15:00	One dose
8	15:30	MSM water ONLY
9	16:00	One dose
9	16:30	MSM water ONLY
10	17:00	One dose
10	17:30	MSM water ONLY
11	18:00	One dose
11	18:30	MSM water ONLY
12	19:00	One dose
12	19:30	MSM water ONLY
13	20:00	One dose
13	TREATMENT COMPLETE	MSM water NOT REQUIRED

It doesn't matter what time of the day you take the actual treatment. That is up to you, but from beginning to end it is 12 complete hours.

As mentioned above, the intent of this treatment is to kill every MICROBE, inside of every cancer cell. Any cancer cell which has all of its microbes killed will, in about a week, revert into a normal cell. As a final comment, this treatment may cause bad breath on the day of the OCC and the day after the OCC.

What To Eat For The Treatment

Pay close attention to the "cancer diet" while on any cancer treatment. AVOID sugar, refined flour, meat (unless the patient is extremely frail), dairy products (except for the Budwig Diet), etc. If you eat whole, raw foods, you will be OK. During this treatment, the wrong foods will feed cancer cells and will interfere with the ability of DMSO to penetrate the cell membranes.

What To Expect From This Treatment

The objective of this treatment is to revert cancer cells into normal cells.

If this is what happens, this is what you can expect from this treatment:

First, reverting cancer cells into normal cells WILL NOT shrink tumors. If the cancer cells are removed from the tumor by this treatment, the body should eventually get rid of any tumors, but it will not happen quickly.

Second, for the same reasons, the OCC will not get rid of any fibrin. What happens to the fibrin when the cancer is gone is unknown at this time.

Third, the OCC SHOULD reduce the pain of cancer within 2 or 3 weeks. Whether the pain is caused by lactic acid or some other cause, most types of pain will be reduced by this treatment. Pain caused by tumors pressing against some other part of the body will not be immediately affected by this treatment.

Fourth, a cancer patient who has very low energy levels can expect an increase in energy within a couple of weeks. As cancer cells are reverted into normal cells the non-cancerous cells get a boost of energy because the cancer cells are no longer stealing glucose and nutrients from the non-cancerous cells.

Fifth, the OCC should stop the spread of cancer. This will not be obvious for awhile; but without any cancer cells, there is no reason for the cancer to spread. This, of course, is more important for fast-spreading cancers.

Sixth, swelling and inflammation should also be reduced by this treatment within a few weeks. However, this will depend on what is causing the swelling and inflammation.

In summary, tumors and fibrin will not immediately be affected by this treatment. These will take the longest to be affected by this treatment. The reduction of fibrin may be helped by taking proteolytic enzyme supplements, also known as pancreatic enzyme supplements.

How You Can Help Other Cancer Patients!!

This treatment can never be perfected without information from people who have used the treatment. It is this information, and nothing else, which will allow us to fine-tune this treatment for other cancer patients. If you do not contact us, then you are only benefiting yourself, not others, with information that may help them. REMEMBER that it will take about two weeks to notice ANY difference because it will take about two weeks for the cancer cells to make the adjustment in their metabolism to become normal cells again.

The second most important thing you can do for other cancer patients is keep a daily diary of your experiences on this treatment, whether you take the OCC once or twice or three times. The main things to include in the diary are how you feel, any unexpected symptoms, a general description of your diet (not too detailed), and similar types of information.

Start the diary the day before the OCC and explain exactly how the cancer patient (which may be you or someone else) feels. Then keep the daily diary going for at least three or four weeks after the OCC, explaining how the cancer patient feels every day. Also include the results and dates of CT scans or PET scans.

Email this accumulating diary to the damaansax@vodamail.co.za every two weeks!!

Contact DAMAANSA HOLDINGS whether you have questions or comments or need to tell us you are taking the protocol, email support will be provided. Be sure to include "OCC" OR faith drops somewhere in your Subject line so your email is not mistaken for spam. See the contact details on the last page of this manual.

TABLE OF RECOMMENDED DOSAGES

GENERAL EXPLANATIONS

**	Wherever these asterisks appear – it is recommended that the protocols are used in conjunction with each other
Grey Shaded Area	All Protocols that are suggested as a combined treatment are shaded in grey, and we recommend that they be used in conjunction with each other
Orally	Always mix the recommended drops with an equal number of activator together in a glass, wait the prescribed 3 minutes, add a mixer as per Chapter 7, and drink immediately.
Transdermally	Always mix the recommended drops with an equal number of activator together in a glass, wait the prescribed 3 minutes, add the prescribed amount of DMSO, wait an additional 3 minutes – then apply to the skin using your bare hands. NEVER use latex gloves. In the event of risk of contamination to the person applying the DMSO – (in the case of a patient with open sores from AIDS/HIV – apply to an area that does not present open sores, or use a wooden spatula or the back of a solid silver / stainless steel spoon to apply.
Intravenously	Always add the recommended number of drops of FAITH, UNACTIVATED to a 500cc saline solution drip. Wait 1 hour and then follow normal protocols for inserting an intravenous drip. Entry ratio is 1 x 500cc : 1-2 hours
Anally	Always mix the recommended drops with an equal number of drops of ACTIVATOR – wait the prescribed 3 minutes then mix ACTIVATED solution with 30cc of water. Using a syringe and soft tube – insert into the anus. Alternatively use an enema pump.
Allopathic Medicines	It is quite safe to combine FAITH Drops with other medicines EXCEPT other alternative medicines. We prefer that you use ONLY one protocol at a time.
DMSO	Dimethyl Sulfoxide. DO NOT use in conjunction with cortisone medication. Wait 8 hours before starting DMSO protocol.
Atropine	DO NOT use in conjunction with DMSO protocol
Adrenaline	DO NOT use in conjunction with DMSO protocol

**** INDICATES THAT PROTOCOLS SHOULD BE COMBINED**

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Abscesses / boils - Orally	3	3	Day 1 to 7 4 x Daily thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Abscessed teeth – Orally	6	6	2 x Daily – until clear	Mix activated drops with 125ml (125cc) water then rinse out the mouth. Do not leave this solution in the mouth for longer than 60 seconds. In our experience, the pain of an abscessed tooth has been overcome by the immune system in about four hours and patient presents a healthy mouth in less than three weeks. EXCEPTIONS: An abscessed tooth where the abscess is inside the tooth and the FAITH mixture cannot reach it through a hole or some other way, the FAITH treatment will not then handle the abscess.
Aids – Orally – Initial & Maintenance Dose **	3	3	2 x Daily – to maintain	FAITH does not attack AIDS, the immune system does. FAITH Drops provide the immune system with the ammunition. Some cases may only clear up with intravenous injections which may be administered only by a doctor. Start out with the Initial Oral dose and then combine that with the Intravenous, Transdermal and Anal Protocols ALL AT THE SAME TIME . Thereafter continue with Maintenance Dose. Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Aids - Intravenously **	Day 1 to 10 2 x Daily	Calculate 1 drop per 10kg of body weight. Add UNACTIVATED Drops to 500ml (500cc) drip.
Aids – Transdermally DMSO Protocol **	20	20	Day 1 to 20 2 x daily	Mix drops and activate – wait the prescribed 3 minutes and then mix with 2.5ml (2.5cc) (1/2 a teaspoon) DMSO – then wait a further 3 minutes. Thereafter, add 1.25ml (1.25cc) (1/4 of a teaspoon) of water, swirl the solution in the glass to mix well then apply activated drops topically using your bare hands. DO NOT USE LATEX GLOVES TO APPLY
Aids – Anally **	3	3	Day 1- 20 3 x Daily	Mix drops and activate, wait the prescribed 3 minutes and then mix with 30ml (30cc) water.
Allergies - Orally	3	3	2 x Daily – until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
A.L.S. – Lou Gehrig Disease - Orally	3	3	3 x Daily - until tests clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Anaemia – Orally	3	3	Day 1 to 7 4 x Daily thereafter 3 x Daily until normalised	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Arteriosclerosis – Orally	3	3	3 x Daily Until tests clear	As you are aware there are many causes for hardening and clogging of the arteries, but cholesterol seems to be one of the best known reasons. Several people have reported to date that taking the FAITH has reduced the closing of the arteries due to cholesterol. One example was a patient that was told that her veins were 80% clogged. Improvement was documented after FAITH treatment
Arthritis - Orally	3	3	Day 1 to 7 4 x Daily thereafter 2 x Daily - until clear	Ascertain the type of arthritis being treated as FAITH will help with rheumatoid arthritis and Lyme arthritis and some others, but not the tissue damage with normal arthritis.
Arthritis - Lyme - Orally	3	3	2 x Daily – until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers. This is a long term treatment
Asthma - Orally	3	3	2 x Daily – until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers. This is a long term treatment
Athlete's feet - Topically	10	10	4 x Daily – until clear	Mix activated drops with 125ml (125cc) of water. Apply the solution to affected area. Wash off after 3 minutes and dry well.
Bacteraemia (Blood-poisoning) - Orally	3	3	4 x Daily – until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Bedsores Diabetes sores / ulcers	10	10	Every 2 hours until healing begins	Mix activated drops with 60ml (60cc) of water and pour into a spray bottle. Spray bed sore with a fresh solution every 2 hours until healing takes place.
Bronchitis - Orally	3	3	4 x Daily - until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Bites - animal, insect, human, snake or spider - Orally	3	3	Hourly for 4 hours thereafter 2 hourly until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Cancer - Orally * *	2	2	Day 1 to 7 4 x Daily thereafter 3 x Daily until tests are clear	Chlorine dioxide gives the immune system almost perfect ammunition to attack cancer cells. We believe the theory has proven true as many people with cancer have reported that the cancer went away or reduced greatly after taking FAITH. The results we have received have proven to be better results than that of standard medical treatment. We have observed that skin cancer usually dries up and drops off within one week of taking FAITH three times a day. We do not claim that FAITH caused the effect, but we did observe the effect. It is our opinion that when one is working with cancer and it is not getting better an over ruling concept is that you are not taking or administering enough FAITH drops. Ideally we prefer the "AGGRESSIVE MIX" but the dosage alongside could be used to start the person off. If this does not clear the problem - then make use of the "Aggressive Mix" dosage. CONTINUE VISITING YOUR DOCTOR AND TAKING ANY PRESCRIPTION MEDICINES. MAKE SURE YOU CONTINUE WITH ANY RADIATION OR CHEMOTHERAPY PRESCRIBED FOR YOU. Use all 4 protocols below - together
Cancer - Transdermally - DMSO - (OCC Protocol) recommended * *	20	20	<u>Day 1</u> 1 x per Hour for 4 hours <u>Day 2</u> 1 x per Hour for 13 hours <u>Day 3</u> Rest. <u>Day 4</u> 1 x per Hour for 4 hours <u>Day 5</u> 1 x per Hour for 13 hours <u>Day 6</u> Rest <u>Day 7</u> 1 x per Hour for 4 hours <u>Day 8</u> 1 x per Hour for 13 hours	Mix drops and activate – wait the prescribed 3 minutes and then mix with 2.5ml (2.5cc) (1/2 a teaspoon) DMSO – then wait a further 3 minutes. Thereafter, add 1.25ml (1.25cc) (1/4 of a teaspoon) of water - swirl to mix well then apply topically using your bare hands. DO NOT USE LATEX GLOVES TO APPLY See OCC Protocol in Chapter 8 of Faith Protocols

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Cancer - Intravenously * *	1	Nil	Day 1 to 10 2 x Daily	Calculate 1 drop per 10kg of body weight. Add UNACTIVATED drops to a 500ml drip, twice daily.
Cancer – Anally * *	1	1	Day 1 to 10 3 x Daily	Calculate 1 drop per 10kg of body weight. ACTIVATE , wait prescribed 3 minutes and mix with 30cc of water.
Candida – Thrush - Orally	3	3	Day 1 - Hourly thereafter	Mix activated drops with 125ml (125cc) water and rinse out the mouth. Do not leave this solution in the mouth for longer than 60 seconds
			4 x Daily - until clear.	
Candida – Thrush – Topically	10	10	3 x Daily - until clear.	Mix the activated drops with 20ml (20cc) of water. Apply the solution to affected area.
Cellulites - Orally	3	3	4 x Daily - until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Cholesterol - Orally	3	3	2 x Daily – until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Chicken-pox – Orally	2	2	Day 1 – Hourly thereafter	Mix activated drops with 75ml of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			4 x Daily - until clear	
Cholera - Orally	3	3	Day 1 – Hourly	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Day 2 - Every 2 hours not more than 4 x Daily	
			Day 3 - Every 4 hours - not more than 2 x Daily	
Common Cold - Orally	3	3	4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Crabs – lice – Topically	10	10	Repeat every 2 hours until clear.	Dilute activated drops into 125ml (125cc) of water. Apply to area and rinse off after 3 minutes
Croup – Orally	2	2	4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Cystitis – Orally	3	3	Day 1 - 4 x Daily thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			3 x Daily until clear.	
Cysts – Orally	3	3	Day 1 - 4 x Daily thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			2 x Daily until clear	
Dandruff – Topically	10	10	2 x Daily until clear.	Dilute activated drops into 125ml (125cc) of water. Apply to area and rinse off after 3 minutes.
Dermatitis - Orally	3	3	4 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Diabetes - Orally	2	2	Day 1 – Hourly thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			4 x Daily until normalised	
Diarrhoea - Orally	3	3	Day 1 – Hourly thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Day 2 – Every 2 hours not more than 4 x Daily	
			Day 3 - Every 4 hours not more than 2 x Daily	
Diverticulitis - Orally	3	3	Day 1 - 4 x Daily thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			3 x Daily - until clear.	
Eczema - Orally	3	3	3 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Encephalitis - Orally	3	3	Day 1 - hourly thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			4 x Daily until clear.	

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Eye Infections - Stye Conjunctivitis - Iritis - Orally * *	3	3	Day 1 - hourly thereafter Day 2 - Every 2 hours Day 3 – Every 2 hours Thereafter take Maintenance Dose of 3 drops 3 x Daily for ten days	Combine with Eye Rinse Protocol as below
Eye Infections - Stye Conjunctivitis Iritis Extreme Eye Infections Eye Rinse Protocol * *	1	1	Repeat every 2 hours until infection has cleared.	Combine with Oral Treatment as above FOLLOW INSTRUCTIONS WITH EXTREME CARE, AS YOU CAN CAUSE DAMAGE TO THE EYES. <ul style="list-style-type: none"> ○ Use one drop of FAITH in 15ml (15cc) (1 tablespoon) of distilled water or eye drops. ○ Wash eye with this solution, applying directly into the eye using an eye rinse bowl. ○ Allow it to remain in the eye for 15 seconds. ○ SET A TIMER. DO NOT GUESS AT THE TIME. ○ Then use distilled water or eye drops to wash the eye until you are certain that you have washed every bit of liquid out of the eye. Rather over rinse than under rinse. ○ Repeat this every two hours until the infection has cleared. ○ If you are unable or unwilling to administer the drops directly into the eyes, you can depend upon the doses by mouth if it is done every hour. ○ REMEMBER: DO NOT allow even one drop of FAITH mixture to remain in the eye, If you do it will cause a burn. ○ In one case where a person did not rinse their eye sufficiently, assuming that the solution was so ineffectual that it would not cause any harm; the result was the skin surrounding the eye drooped considerably and it appeared as if the eye would actually fall out of the socket. This however did clear up after a few days. But, are you prepared to take the risk of it not clearing up. ○ So, pay attention and don't guess the formula mix or the time you allow it to sit in the eye nor take for granted the fact that you must thoroughly flush the eye after 15 seconds.
Flu – Orally	3	3	Day 1 - 4 x Daily thereafter 3 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Food poisoning – Orally	3	3	Hourly for 4 hours thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Every two hours until clear	
Gastritis - Orally	2	2	Day 1 - 4 x Daily thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			3 x Daily until clear	
Glandular fever – Orally	3	3	Day 1 - 4 x Daily thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			3 x Daily until clear.	
Gout – Orally	3	3	Day 1 to Day 7 – 4 x Daily thereafter 2 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Head Sores – Topically	10	10	1 x daily - repeat until clear	Add 125ml (125cc) water to the 10-drop ACTIVATED solution of FAITH. <ul style="list-style-type: none"> ○ When combed into the hair with enough solution to penetrate to the scalp it will cure most head sores in a few hours. ○ If you use it daily it makes for a healthy scalp, but it also opens the cuticle scale of the hair and strips the natural melanin out of the hair. So, if you are not a natural blonde we suggest you use it once or twice a month. ○ When used daily you begin to notice that your is bleaching blond after a few days. (It is a very nice blond.)
Heart - Pericarditis Orally	1	1	Day 1 – Hourly thereafter	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			2 drops every 2 hours until clear.	
Hepatitis A – Orally	2	2	Day 1 - Every hour	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Day 2 – 3 drops every 2 hrs	
			Day 3 - 3 drops 4 x Daily until clear.	

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Hepatitis B – Orally	2	2	Day 1 – every hour	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Day 2 – 3 drops every 2 hours	
			Day 3 – 3 drops 4 x Daily until clear.	
Hepatitis C - Orally	2	2	Day 1 – 2 drops every hours	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Day 2 – 3 drops 4 x daily until clear.	
Herpes – Orally	3	3	4 x Daily	Mix activated drops with 75ml (75cc) of water and drink. Herpes virus should be gone in about two weeks. However, in very bad cases, it might take up to two months to finally overcome the Herpes virus. With severe cases take the formula every two hours for several days. This aids the FAITH drops to penetrate deeper into the tissues. In worse cases, one may have to use the drops intravenously, by calculating 1 drop per 10kg of body weight. Add UNACTIVATED drops to 500ml drip twice daily. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Infected Gums – Orally	6	6	2 x Daily	Mix activated drops with 125ml (125cc) of water and use this solution to rinse the mouth. Do not leave this solution in the mouth for longer than 60 seconds.
Insect Bites – Topically	10	10	As required	Although this is not particularly dangerous or poisonous - more irritating, yet a severe mosquito bite can be very uncomfortable for several hours and if continually scratched can become infected. Use this solution to apply dermally by gently tapping the solution onto the bites with your fingertips to help the solution to penetrate. The itching from the bite will be gone in 5 minutes with the bump showing signs of diminishing almost immediately. You do not need to wash this solution off, for once you have added the activator to the drops you have negated its ability to burn your skin. Once citric acid has been added the alkalinity of the solution has been destroyed. However, when using the solution in larger areas of the body than a small insect bite always add about 75-125ml (75-125cc) of water to the 10-drops solution that you mix - depending on the area needing to be treated.
Jock Itch – Topically	6	6	3 x Daily – until clear.	Mix activated drops with 125ml (125cc) of water. Apply the solution with your finger tips to affected area. Rinse the area after 3 minutes as the area is very sensitive.

	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Kidney- Pylonephritis - Orally	2	2	4 x Daily – until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Laryngitis – Orally	2	2	4 x Daily – until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Leukemia – Orally	3	3	2 x Daily – until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Lupus – Orally	3	3	2 x Daily – until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Lyme Disease – Orally	3	3	Day 1 to 14 Every 2 hours thereafter 3 x Daily - until clear.	Mix activated drops with 75ml (75cc) of water and drink. It might take as long as a year to overcome Lyme Disease. There is no proof yet but the information we have received is that everyone who has Lyme Disease and is taking FAITH is experiencing improvement. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Malaria - Orally	10	10	Day 1 – 2 x hourly thereafter 4 hourly until tests negative or symptoms have cleared.	Mix activated drops with 75ml (75cc) of water and drink. The procedure for malaria is to always start with a 10 drop dose, and give a second 10 drop dose in one to two hours. Most of the symptoms should have cleared within four hours of the second dose. Should the symptoms persist after the second dose, simply give a third dose. All malaria parasites should be dead after second dose, so, if the patient is still sick, it will not be from malaria, but rather from some other disease. In that case, continue with at least two 10 drop doses each day until they are feeling well. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Measles	1	1	Day 1 - 2 x hourly thereafter Every 4 hours thereafter until clear.	Mix activated drops with 75ml (75cc) water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Melanomas – Orally * *	1	1	Begin with 1 drop and increase daily.	Begin the Standard Formula Protocol and increase until you are able to administer the "Aggressive Mix". If necessary, exceed the prescribed number of drops to 15 drops per dose three times a day. We do not normally recommend such high individual doses but as it is usually only about a week or two before you notice the shrinkage. This shrinkage can often be indicated by a white ring around the melanoma and it then it becomes smaller and smaller until after about two weeks when it is usually gone from the surface. You would obviously need to do all necessary tests to confirm this. <u>Feedback we have received is:</u> Small cancerous sores on the body or hands that have not responded to treatment sometimes for over a year, treat with 1 drop of FAITH mixed with 3 drops of the ACTIVATOR , wait 3 minutes and then add 2 drops of DMSO. This sometimes can be quite painful as it burns excessively but the pain usually is alleviated after 3 minutes.
Melanoma – Transdermally * *	5	5	4 x Daily until it falls off	Mix activated drops in a tot measure and wait prescribed 3 minutes. Mix in 5 drops (using an eye dropper) of DMSO into the activated solution. Wait an additional 3 minutes. Mix this solution with 5ml (5cc) of water. Invert tot measure over the melanoma and hold firmly in place for 3 minutes. This will ensure that the melanoma is totally submerged in the FAITH solution.
Meningitis – Orally	2	2	Day 1 – Hourly thereafter 2 hourly thereafter until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Mumps – Orally	2	2	Every 3 hours until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Obesity – Orally	2	2	3 x Daily	Calculate 1 drop for every 10kg of body weight. Increase dose by 1 drop daily until full dose reached according to weight. Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Osteomyelitis – Orally	3	3	Day 1 – 2 hourly thereafter 4 hourly until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Otitis - infections	2	2	4 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Pancreatitis – Orally	2	2	Day 1 to 2 4 hourly thereafter 4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.

LLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Pneumonia – Orally	3	3	Day 1 – 4 hourly	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Thereafter 4 x Daily until clear.	
Pyorrhoea – Orally	6	6	2 x Daily until clear.	Activate drops and use solution to brush the teeth. Make and use a new solution every morning. Do not leave this solution in the mouth for longer than 60 seconds, as it might damage your sight, there are people who regained their sight merely by replacing the metal in their mouth.
Ringworm – Orally * *	3	3	4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Ringworm – Topically * *	10	10	Repeat every 3 hours until clear	Mix activated drops in a tot measure and wait prescribed 3 minutes. Mix in 5 drops (using an eye dropper) of DMSO into the activated solution. Wait an additional 3 minutes. Mix this solution with 5ml (5cc) of water. Invert tot measure over the ring worm and hold firmly in place for 3 minutes. This will ensure that the ringworm is totally submerged in the FAITH solution.
Scabies – Orally	3	3	2 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Scabies – Topically	10	10	Repeat every 3 hours until clear	Dilute with 125ml of water. Apply to area then rinse off after 3 minutes.
Septicaemia – Orally	3	3	Every hour for 1 day thereafter every 4 hours thereafter until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Shingles / Herpes - Orally	2	2	Every 2 hours for 1 day thereafter 3 drops 4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Sinusitis - Orally	3	3	Every 4 hours until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Spider Bites - Brown Recluse Spider – Orally * *	3	3	3 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. Use in conjunction with Topical Protocol below. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Spider Bites - Brown Recluse Spider – Topically * *	10	10	3 x Daily for two days	Mix activated drops in a tot measure and wait prescribed 3 minutes. Mix in 5 drops (using an eye dropper) of DMSO into the activated solution. Wait an additional 3 minutes. Mix this solution with 5ml (5cc) of water. Invert tot measure over the spider bite and hold firmly in place for 3 minutes. This will ensure that the bite is totally submerged in the FAITH solution. In conjunction with this apply any nappy rash ointment containing at least 40% zinc oxide, 5 minutes after the application of the DMSO protocol. Nappy rash ointment containing at least (higher if possible) 40% zinc oxide will kill most of the poison and reduces the pain and itching of a brown recluse spider bite in less than one day. We have seen many people treat this bite successful with this nappy rash ointment.
Spider Bites - Black Widow – Orally * *	3	3	3 x Daily for 2 to 3 days to aid the body in recovering	Mix activated drops with 75ml (75cc) of water and drink. Use in conjunction with Topical & Intravenous Protocols below. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Spider Bites - Black Widow – Topically * *	10	10	3 x Daily for 2 to 3 days to aid the body in recovering	Mix activated drops in a tot measure and wait prescribed 3 minutes. Mix in 5 drops (using an eye dropper) of DMSO into the activated solution. Wait an additional 3 minutes. Mix this solution with 5ml (5cc) of water. Invert tot measure over the spider bite and hold firmly in place for 3 minutes. This will ensure that the bite is totally submerged in the FAITH solution.
Spider Bites - Black Widow – Intravenously * *	3	Nil	3 x Daily for 2 to 3 days to aid the body in recovering	Mix UNACTIVATED drops with 20cc (20ml) of sterile water in a glass. Using a syringe siphon up un-activated solution and inject intravenously, very slowly. (30 seconds) Use in conjunction with Topical Protocol above.
Tuberculosis - Orally	2	2	Day 1 - 2 drops every hours	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
			Day 2 - 2 drops every 2 hours	
			Day 3 - 3 drops 4 x Daily until clear	

ILLNESS	FAITH DROPS	ACTIVATOR	DROPS SCHEDULE	COMMENTS
Tumors – Orally * *	3	3	3 x Daily until tumour has gone. mixers.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable
Tumours – Transdermally * *	20	20	2 x Daily until tests are clear	Mix drops and activate – wait the prescribed 3 minutes and then mix with 2.5ml (2.5cc) (1/2 a teaspoon) DMSO – then wait a further 3 minutes. Thereafter, add 1.25ml (1.25cc) (1/4 of a teaspoon) of water - swirl to mix well then apply topically using your bare hands. DO NOT USE LATEX GLOVES TO APPLY See OCC Protocol in Chapter 8 of Faith Protocols
Urinary tract infections Orally	3	3	4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Ulcers – Orally	3	3	4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Vaginal infections – Orally	3	3	4 x Daily until clear.	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Vaginitis – Orally	3	3	4 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Vaginosis – Orally	3	3	4 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Warts – Topically	10	10	Repeat daily until wart falls off	Mix the drops and apply directly onto wart. Rinse off after 3 minutes.
Weight Control	3	3	3 x Daily until normalised	Mix activated drops with 75m (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Whooping Cough – Orally	2	2	4 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.
Yeast infection – orally	3	3	3 x Daily until clear	Mix activated drops with 75ml (75cc) of water and drink. See Chapter 7 – Page 53 of Protocols for various acceptable mixers.